

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C12M 3/04, 3/10, C12Q 1/02, 1/24	A1	(11) International Publication Number: WO 95/29983 (43) International Publication Date: 9 November 1995 (09.11.95)
(21) International Application Number: PCT/US95/05117 (22) International Filing Date: 26 April 1995 (26.04.95) (30) Priority Data: 08/234,448 28 April 1994 (28.04.94) US (71) Applicant: THE PENN STATE RESEARCH FOUNDATION [US/US]; 207 Old Main, University Park, PA 16802 (US). (72) Inventors: HAMMERSTEDT, Roy, H.; 1119 Kathryn Street, Boalsburg, PA 16827 (US). BARBATO, Guy, F.; 651 West Hillside Avenue, State College, PA 16803 (US). CRAMER, Palmer, G.; 102 Wheel Circle, State College, PA 16803 (US). (74) Agent: MONAHAN, Thomas, J.; The Pennsylvania State University, Intellectual Property Office, 113 Technology Center, University Park, PA 16802 (US).		(81) Designated States: CA, JP, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE). Published <i>With international search report.</i>
(54) Title: PROCESS FOR EVALUATING FERTILITY OF SPERMATOOZOA SAMPLES (57) Abstract A method for testing the potential fertility of spermatozoa in a sample by contacting an aliquot of the sperm sample onto a protein extracted from native vitelline membranes removed from chicken or turkey eggs to enable quantification of the number of sperm which bind to the protein. Typically, the vitelline membrane protein is coated onto a solid substrate and the protein extract is first prepared by dissecting vitelline membranes (lamina perivitellina plus lamina extravitellina) from either a group of chicken eggs or a group of turkey eggs, rinsing the membranes free of albumin and yolk, and subdividing the membranes into small particles. The resulting protein is heat solubilized, purified with centrifugation and concentrated to yield the protein extract in solution. The extracted protein then is used as a protein substrate for spermatozoa binding, with sequential contacting of the protein extract with the spermatozoa and measuring the amount of sperm-egg binding that takes place. Intact egg membranes may alternatively be used to bind sperm. Correlation between sperm-egg binding and fertility of the spermatozoa in the sample is direct and linear.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	GB	United Kingdom	MR	Mauritania
AU	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgyzstan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LI	Liechtenstein	SK	Slovakia
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CN	China	LU	Luxembourg	TD	Chad
CS	Czechoslovakia	LV	Latvia	TG	Togo
CZ	Czech Republic	MC	Monaco	TJ	Tajikistan
DE	Germany	MD	Republic of Moldova	TT	Trinidad and Tobago
DK	Denmark	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	US	United States of America
FI	Finland	MN	Mongolia	UZ	Uzbekistan
FR	France			VN	Viet Nam
GA	Gabon				

PROCESS FOR EVALUATING FERTILITY OF SPERMATOOZOA SAMPLES

Field of the Invention

The present invention relates to an evaluation process for assessing the fertility of avian and mammalian spermatozoa.

5

Background of the Invention

Assisted reproductive technologies such as artificial insemination have been practiced commercially with livestock and horses for almost a century, and with poultry, humans and other species for almost a half century. Artificial insemination requires a sample of semen consisting of spermatozoa together with seminal plasma contributed by the epithelial cells of the epididymis and accessory organs. In addition to water, seminal plasma contains numerous proteins and glycoproteins, phospholipids, lipids, carbohydrates and ions.

It was recognized early on that a sample of semen contains a heterogeneous population of sperm, with some having--and others lacking--fertilizing potential. Thus, potential fertility differs from sample to sample for a given male, and differs greatly among different males of a species. Prediction of potential fertility is a recurring question in human medicine and in research on sperm from animals of economic importance. A potential effect on fertility is intrinsic to any attempt to improve methods for preservation of sperm, or in efforts to solve the many unanswered mysteries of sperm biology. Unfortunately, valid assessment of fertility of an individual male has heretofore been expensive for species such as cattle and chickens, impractical and expensive for horses, and impossible for humans.

For many years it has been recognized that potential fertility of a given seminal sample is difficult to measure accurately. Numerous approaches have been used, but no single method tests all of the attributes of sperm function that contribute to fertility. Tests evaluating percentage of motile sperm, motion characteristics of sperm, or exposed carbohydrates or membrane integrity are somewhat useful, but are

not reliably predictive. Motile sperm with a normal morphology are more likely to be fertile than immotile sperm or sperm with a bizarre shape or structure. Nevertheless, neither subjective nor objective measurements of sperm motility or morphology have proven to be highly predictive of potential fertility.

5 A review of the binding mechanism between spermatozoa and eggs is instructive in this context. The initial event in the fertilization process is the binding of one or more sperm to the egg investments. In mammals, a spermatozoon is considered to bind to the oocyte through a series of egg-binding proteins, collectively called ligands, located on the surface of the spermatozoon which interact with
10 appropriate members of a series of sperm receptors located on the investments of the oocyte, namely, the zona pellucida and oocyte plasma membrane. This terminology for location of binding proteins on the sperm and receptors on the egg is most logical, is traditional with invertebrates and applicable to mammals, and will be used hereafter with reference to both mammals and birds.

15 Egg-binding proteins of mammalian sperm generally are considered to be transmembrane proteins or glycoproteins, possibly with enzymatic activity, with an extracellular domain that interacts with a specific sperm receptor. The consensus scenario which has evolved is one of sequential binding: first a loose binding to the zona pellucida via one or more molecule(s) located on the sperm plasma membrane;
20 then a tight binding of the sperm to the zona pellucida via molecule(s) located on the plasma membrane and/or the acrosomal membrane; followed by tight binding of the sperm to the oocyte plasma membrane followed by fusion of the sperm and egg plasma membranes and entrance of the entire spermatozoon into the oocyte. Thus, species-specific adhesion between sperm and eggs is attributable to complexes formed between
25 egg-binding proteins on sperm and complementary sperm receptors on eggs. At least one species non-specific egg-binding protein also must be located on the sperm surface.

 Eggs also have receptors for sperm ligands. In mammals, the plasma membrane of the egg is overlain by the zona pellucida, an acellular coating formed around the oocyte while it is still within the ovarian follicle. Depending on the species,
30 the zona pellucida may limit access of supernumerary sperm to the plasma membrane. In common mammals, fertilization apparently follows a similar process which features

a pivotal role for the highly conserved zona pellucida, comprised of several glycoproteins in the mouse termed ZP₁, ZP₂, and ZP₃. A loose non-specific binding of a spermatozoon to the zona pellucida is followed by strong attachment of sperm ligands to receptors on ZP₃, initiation of the acrosome reaction and conversion of proacrosin to acrosin (a serine protease), binding to ZP₂, enzymatic digestion of a channel for that spermatozoon through the zona pellucida and passage of the spermatozoon into the underlying perivitelline space, after which other sperm ligands bind to receptors on the oocyte plasma membrane. It is unclear if sperm motility is requisite for these binding responses, but motility certainly is necessary for penetration of the zona pellucida. Obligatory steps include (1) initial loose binding of a spermatozoon to the zona pellucida and (2) subsequent tight binding by receptors on ZP₃. Given the pivotal role in fertilization of penetration of sperm through the zona pellucida, it is not surprising that the number of sperm trapped in the zona (termed accessory sperm) is related to the fertilization status of the oocyte therein; the more sperm, the greater the probability that fertilization occurred.

In birds, the maternal gamete and yolk of an ovulated egg is encircled by a multilayered egg membrane (vitelline membrane), comprised of a distinct inner lamina perivitelline and an outer mesh-like lamina extravitellina. After passage of the egg from the infundibulum into the magnum of the oviduct, a series of albumin coats are added followed by components associated with the shell. Avian sperm normally can penetrate the lamina extravitellina and bind to the lamina perivitelline, steps essential for fertilization, only during the approximately 15 minutes which elapse between ovulation and deposition of the initial albumin coat. Accessory sperm can be found on or within the egg membrane of a fertilized egg. Little research has been done on the molecular nature of the egg membranes or sperm-egg binding in chickens. However, it is known that the lamina perivitelline is composed primarily of 3 glycoproteins while the lamina extravitellina is a mixture of other components.

There have been to date countless approaches to evaluate sperm quality, and correlation of test results with known fertility data is not the same as accurately predicting fertility in an unknown sample--which has not been done to date. Accordingly, a need remains for a simple and accurate test to assess the fertilization

potential of spermatozoa in a single sample from an individual mammalian or avian organism. Such a test is useful both in human health care and commercial animal breeding.

Summary of the Invention

5 The present invention is a method for testing the potential fertilizing capability of spermatozoa by placing a suspension of these spermatozoa onto a protein extracted from native vitelline membranes removed from chicken or turkey eggs, with the protein usually having first been coated onto a solid substrate, to enable quantification of the number of sperm which bind to the protein. Typically, the
10 vitelline membrane protein extract is prepared by dissecting vitelline membranes (lamina perivitelline plus lamina extravitellina) from either a group of chicken eggs or a group of turkey eggs, rinsing the membranes free of albumin and yolk and subdividing the membranes into small particles mechanically by ultrasonic energy shearing. The resultant divided protein is solubilized, purified, centrifuged and
15 concentrated to yield the protein extract in solution. The extracted protein then is used as a protein substrate for spermatozoa binding, with sequential contacting of the protein extract with the spermatozoa and measuring of the amount of sperm-egg binding that takes place. Alternatively, the washed egg membrane, before shearing and solubilization, can be laid on a solid substrate to provide the binding surface.
20 Correlation between sperm-egg binding and fertilizing potential of the spermatozoa is direct and linear.

Brief Description of the Figures

Figure 1 is a perspective view of a microscope slide, coated with an intact membrane according to the invention, covered by a template with holes to form
25 wells in which to assay sperm fertility.

Figure 2 is a graph showing the results of a bull sperm binding test described in Example 10, below.

Figure 3 is a bar graph showing the results of a horse sperm binding test described in Example 11, below.

Figure 4 is a graph showing the results of a mouse sperm binding test described in Example 12, below.

5 Figure 5 is a set of three graphs showing the results of human sperm binding tests described in Example 13, below.

Detailed Description of the Invention

The present invention is a method for testing the fertility of a sample of spermatozoa by contacting aliquots of the sample onto a protein extracted from native
10 vitelline membranes removed from chicken or turkey eggs, or onto the vitelline membranes themselves, to enable quantification of the number of sperm which bind to the protein. Typically, when the protein extract method is used, the vitelline membrane protein extract is prepared by dissecting vitelline membranes (lamina perivitelline plus lamina extravitellina) from either a group of chicken eggs or a group
15 of turkey eggs, rinsing the membranes free of albumin and yolk and subdividing the membranes into small particles by applying mechanical ultrasonic shearing. The resultant protein then is solubilized in buffer with heat, purified with centrifugation and concentrated (for example with a 10 kDa cutoff ultrafiltration membrane) to yield the protein extract in buffered solution. The extracted protein then is used as a protein
20 substrate for spermatozoa binding, with sequential contacting of the protein extract with the spermatozoa and determining the number of sperm bound because of sperm-egg binding reactions. The determination can be visual counting by microscopy, or an automated equivalent using commercially available microscopic image analysis software, or the use of commercially available microwell plate readers quantifying
25 absorbance or fluorescence or stained spermatozoa. Correlation between sperm-egg binding and fertility (fertilizing ability) of the spermatozoa in the sample is direct and linear.

More particularly, the present method is conducted as follows.

The vitelline membranes are dissected from chicken or turkey eggs (one or the other but not both, either fresh unfertilized or stored up to 6 months at 2-5° C) and rinsed free of albumin and yolk by repeated immersion and shaking in repeated washes of cleansing buffer. A typical cleansing buffer contains 50 mM sodium phosphate and 145 mM sodium chloride, pH 7.2, but equivalent buffers can be substituted. In general, throughout this specification any buffer may be substituted for any other buffer as long as the buffer is generally acceptable for suspending or washing biological materials, according to ordinary skill in the art. The washed vitelline membranes are either used immediately or are stored in cleansing buffer at 2-5° C for up to several days before use. The membranes then are subdivided into small particles by applying mechanical or ultrasound energy. The resultant material is solubilized, purified with centrifugation and concentrated to yield an egg membrane protein extract in solution.

By "protein," in the context of the present invention and the protein found in the egg membrane which is capable of binding spermatozoa, applicant means any biomolecule containing an amino acid chain or peptide moiety including but not limited to lipoproteins and glycoproteins. In other words, for the purpose of this invention a broad definition of protein has been adopted and is intended.

The suspensions are pooled and the total volume brought to 6 mL/membrane with additional buffer. The suspensions are heated in a closed vessel at 73° C for 20 minutes to solubilize the protein, cooled to 4° C, and centrifuged at 27,000 x G at 4° C for 15 minutes. The supernatant protein solution is aspirated and concentrated using a 10-kDa cutoff ultrafiltration membrane to provide a solution containing approximately 2.5 mg/mL protein. Other methods to extract the protein are possible, and the active protein molecule(s) can be purified by any method. The solution can be used immediately to coat slides or microwell plate wells, or stored at -20° C for at least 3 months.

Subsequently, the cavities in a microwell plate are coated with the protein extract from vitelline membrane (a concentration of 2.5 mg/mL protein in 20 mM Tris-HCl buffer, pH 8.0, is typical of one series of extracts used) to obtain a thin,

uniform layer. With purer membrane extracts, less material need be coated to the substrate. The protein should be rapidly dried, using low heat applied under vacuum, although the protein can be air dried. After the coating is dry, the coated plate can be used immediately or stored in a humidified cold environment.

5 In order to use the microscope slide prepared as above, a template is laid over a clean intact egg membrane placed on a microscope slide. The template is simply a generally flat construct with holes in it so that upon overlaying the template onto the egg membrane, individual reaction wells result. The template may be made of any material, such as silicone rubber, which does not bind spermatozoa and is
10 nontoxic to sperm. The template can be bonded to the slide with nontoxic adhesive (e.g., UV-catalyzed adhesives known in the art) or clamped in place, and the slide/template combination can be used immediately or stored under humidified cold conditions.

Referring now to Figure 1, the combined prepared egg membrane,
15 microscope slide and template are shown as the well device 10, having the microscope slide 12 having an egg membrane 14 (or protein coating) thereon, with the microscope slide 12 being bonded with a peripheral layer of bonding agent (not visible) to a template 16 having holes 18 therein. The invention embraces any well devices having the present egg membrane or protein coating deposited therein, so Figure 1 is merely
20 exemplary.

For use, the wells in the device are rinsed with buffer and used to assay for sperm utility, as follows.

Spermatozoa in suspension with autologous seminal plasma and added buffer, or preliminarily washed free of seminal plasma, are added dropwise to the wells
25 formed by the template and the microscope slide. Incubation is conducted by leaving the sperm in the well for about 1-3 hours at 37° C in a humidity chamber. After incubation, the sperm suspension is decanted, and each well is rinsed with buffer, with agitation, for 3 minutes. A method for consistent washing of membranes is crucial for reproducible results.

After rinsing, the bound sperm are stained with 4,6-diamidino-2-phenylindole (DAPI; 1 μ g/mL in PBS buffer) and are examined under a fluorescent microscope. The number of sperm bound per unit area of membrane is then counted. The correlation between number of bound sperm counted and fertility of the sample is direct and linear.

The present method can be conducted with washed intact chicken egg or turkey egg membranes, if desired, rather than preparing the membranes in solution and coating a solid substrate. The intact membranes are more difficult to handle than the protein extract, however, because of their fragility, the difficulty of arranging them without bubbles or wrinkles on a solid support, and the more difficult viewing of their wrinkled surface under microscopy. Despite these disadvantages, however, the use of egg membranes per se has utility, and thus forms a part of the present invention. In fact, inasmuch as the following examples are presented in the order they were conducted as the present invention was developed, the earliest examples recite the use of intact egg membranes.

The following examples are illustrative, and are not intended to limit the invention in any way.

Example 1

Vitelline membranes (lamina perivitelline plus lamina extravitellina) were dissected from chicken eggs (fresh unfertilized chicken eggs were used but eggs stored up to 6 months at 2-5° C could have been substituted), and rinsed free of albumin and yolk by repeated immersion and shaking in a series of beakers containing swirling cleansing buffer (50 mM sodium phosphate, 145 mM sodium chloride; pH 7.2). The washed vitelline membrane was used immediately, although it could have been stored in cleansing buffer at 2-5° C for several days before use. The membrane was not allowed to dry out between removal from the egg and use in the assay.

Washed vitelline membranes from turkey eggs also were prepared in the same manner.

To prepare the assay device, a microscope slide was cleaned and dried for use. A template of silicone rubber that contained a series of holes of uniform size (3-10 mm across) was prepared. Silicone rubber was selected because it is not toxic to sperm, it can be readily cut to contain holes of a uniform and standard size, and can form a seal when placed between adjacent surfaces. The template was coated with a nonstick substance (Fisher Brand Dricote, known in the art, was used) to prevent adherence of cells to the template materials during the assay.

A segment of vitelline membrane was laid over a 24 x 24-40 mm coverslip, taking care to avoid folds or tears. Immediately the silicone-coated template was laid over the membrane, so that the wells in the template were superimposed over the vitelline membrane. The template was clamped over the membrane using plastic clamps, and the holes in the template formed wells for dropwise addition of sperm suspension or control admixtures.

The wells in the template were rinsed with buffer and shaken to dislodge the excess. Then, a measured aliquot of sperm suspension was placed into each well. The sperm suspension included spermatozoa, buffer and autologous seminal plasma. The sperm were left in the well for 1-3 hours at 37° C in a humidity chamber. After incubation, the sperm suspension was decanted, and each well was rinsed with buffer, with agitation, for 3 minutes. A method for consistent washing of membranes was crucial for reproducible results.

After rinsing, the bound sperm were stained with 4,6-diamidino-2-phenylindole (DAP; 1 µg/mL in PBS buffer) and were examined under a fluorescent microscope. The number of sperm bound per unit area of membrane was counted. Appropriate controls were used in parallel wells, and included: sperm rendered immotile, sperm damaged to remove ligands binding to the membrane, or sperm coincubation with an excess of molecules known to block or enhance the binding reaction.

Initial validation of the general assay procedure used slides and templates in which each well held about 20 µL. To establish that the number of sperm bound was a function of the number of motile sperm in the well, a dose response test was conducted; wells contained 0.15 to 10 x 10⁶ fresh sperm (about 80% motile) or 10

x 10⁶ killed sperm (flash frozen to -196° C 3X). After 3 hours of incubation, unbound sperm were washed away, bound cells were stained, and sperm in a 0.5-mm² area were counted at 156X magnification. A dose response curve was obtained.

Example 2

5 Example 1 was repeated except both fertilized and nonfertilized eggs were used, separately, to compare and to evaluate the ability of sperm to bind to membranes recovered from unfertilized or fertilized eggs of similar freshness. It was found that more motile sperm were bound by unfertilized eggs than fertilized eggs, and that the discrimination between live and killed sperm also was greater. This
10 established that vitelline membranes to be used in the present method are preferably taken from unfertilized eggs.

Example 3

 In order to ascertain the relationship of the in vitro results to fertility of the same samples when used for artificial insemination, the following test was
15 conducted. One volume of rooster sperm was extended with three volumes Minnesota-A buffer containing 12% glycerol, was frozen to -196° C, and was carefully thawed. Glycerol was diluted by serial addition of Minnesota-A buffer and sperm were recovered by centrifugation. The sperm pellet was resuspended in Minnesota-A buffer (standard cryopreservation procedure). Freshly collected sperm (no cryopreservation)
20 were diluted in Minnesota-A buffer to serve as a control preparation. Aliquots of 100 x 10⁶ sperm representing each of the 2 treatments were tested using the assay described herein. Fertility of the sperm suspensions also were established using 17-19 hens/group. The results illustrate that the ranking of the 2 samples obtained with the egg membrane binding assay was far more diagnostic of fertilizing capability than
25 either percentage of motile sperm or percentage of sperm with an intact plasma membrane--see Table 1, below.

Table 1

Comparison of laboratory and fertility assays
of fresh and cryopreserved rooster sperm

Treatment	Percent Motile Sperm	Percent Intact Sperm	Relative Sperm-Egg Binding	Mean Fertility (percent)
Diluted sperm (Control-- No Cryopreservation)	95	100	100	86
Standard Cryopreservation	70	73	18	34

Example 4

In order to determine any changes in fertilizing potential of rooster sperm after storage of the sperm at 4° C for 0, 24 or 48 hours the following test was conducted. After storage of rooster sperm for the above-stated periods of time, aliquots of sperm were evaluated for motility and resistance to hypoosmotic shock (both by methods known in the art), and in vitro binding to an egg membrane as described in Example 1. Other aliquots were used to inseminate groups of 60, 60 and 14 hens with 75, 75 or 130 x 10⁶ sperm for sperm stored at 0, 24 or 48 hours, respectively; 356, 350 and 72 eggs were evaluated. Actual egg fertilizations, extent of egg-membrane binding of sperm and the degree of sperm plasma membrane damage were all determined, under microscopy where necessary, and the results appear in Table 2, below.

Table 2
Comparison of rooster spermatozoal characteristics
in laboratory assays and fertility tests

Treatment	Motility		% Intact Sperm Plasma Membranes	Fertility (%)	Relative Sperm-Egg Binding Assay
	% motile ((%) D/min.))	1/velocity $t^{1/2}$			
Zero Time (No Storage)	62	164	91	34	100
Stored at 6° C for 24 Hours	30	97	93	14	60
Stored at 6° C for 48 Hours	31	90	88	4	30

Few of the sperm had a damaged plasma membrane regardless of storage interval (all values <12% damaged), and based on sperm motility the quality of samples stored for 24 or 48 hours were similar, but were inferior to those examined at 0 hours before storage (approximately 50% reduction). The egg membrane binding assay was the only in vitro sperm quality test which detected a difference in potential fertility between sperm stored 24 hours and sperm stored 48 hours: fertility of samples decreased between 0 and 24 hours and further between 24 and 48 hours. The inventive assay is thus illustrated to have utility in predicting fertility of both fresh and stored sperm specimens, and for stored specimens can even detect loss of fertility over time.

Example 5

The following test was conducted to determine the efficacy of the present technique in evaluating the fertility of turkey spermatozoa.

Membranes from chicken hen eggs were prepared for use with fresh (with a high percentage of motile sperm), killed (by cooling to -196° C; no motile sperm), or 24 hours stored turkey sperm. The assay was performed according to Example 1; each well contained 2.5×10^6 turkey sperm in 20 μ L volume. Results were determined in accordance with Example 1 and yielded the following values:

Fresh spermatozoa (expressed in terms of a normalized value of 100%) bound to the egg membranes; approximately 30% of that amount bound when stored spermatozoa were tested for binding to the egg membranes; and only about 3% of the killed spermatozoa bound to the egg membranes. It was thus readily apparent that fresh turkey sperm became bound to the egg membrane, and that such binding was greatly reduced for motile sperm tested after 24 hours storage. Further, fresh and stored sperm were both bound in far greater numbers than killed sperm. The same numeric results were obtained when egg membranes from turkey hens were used in substitution for the chicken eggs, in a parallel test.

Example 6

Preparations containing rooster, bull and ram sperm were incubated with hen's egg membrane as described in Example 1 and then were examined under electron microscopy. Rooster sperm showed close orientation to the outer portion of the perivitelline membrane, whereas bull and ram spermatozoa did not enter the membrane to the same degree, but rather were embedded in an unknown material adhering to the outer membrane of the hen's egg with sperm heads lined up in parallel. This trial illustrated that spermatozoa from different species bind to egg membrane when subjected to the present assay, and also that the nature of the binding may differ by species.

Example 7

We compared ability of an assay using the present protein extract and native egg membrane for their abilities to discriminate between samples of fresh rooster sperm (samples contained a high percentage of motile sperm), intentionally killed sperm (rapidly frozen to -196° C without cryoprotectant--no motile sperm), and sperm cryopreserved by two different conventional procedures.

The protein extract was prepared as follows. The vitelline membranes of chicken eggs were dissected and rinsed free of albumin and yolk by repeated immersion and shaking in repeated washes of 50 mM sodium phosphate and 145 mM sodium chloride, pH 7.2. The washed vitelline membranes were then subdivided into small particles by applying ultrasound energy. The resultant protein was solubilized by heating in a closed vessel at 73° C for 20 minutes, followed by cooling to 4° C, and centrifuging at 27,000 x G at 4° C for 15 minutes. The supernatant protein solution was aspirated and concentrated using a 10-kDa cutoff ultrafiltration membrane to provide a solution containing approximately 2.5 mg/mL protein. The solution was applied (110 µL/well) to a microwell plate and evaporated under a vacuum and heat (60° C).

Spermatozoa in suspension with buffer and traces of autologous seminal plasma were added dropwise to the wells formed by the template and the microscope slide. Incubation was conducted by leaving the sperm in the well for about 2.5 hours at 37° C in a humidity chamber. After incubation, the sperm suspensions were decanted, and each well was rinsed with buffer, with agitation, for 3 minutes. After rinsing, the bound sperm were stained with 4,6-diamidino-2-phenylindole (DAPI; 1 µg/mL in PBS buffer) and were examined under a fluorescent microscope. The number of sperm bound per unit area of membrane was counted.

Direct comparison revealed that equivalent results were obtained with either assay, as shown in Table 3.

Table 3

Percent cells bound by egg membrane vs. protein extract

Sperm Type	Intact Membrane	Protein Extract
Live Sperm	100	100
Cryopreserved Sperm	70	78
Cryopreserved Sperm (Alt. Technique)	30	33
Dead Sperm	9	7

Example 8

Example 7 was repeated except proteins extracted from chicken eggs versus turkey eggs were compared. Both fresh chicken and turkey sperm bound in similar proportions to protein extracts taken from either chicken or turkey eggs, whereas <5% of the killed sperm bound to either extract. Frozen thawed sperm from either species bound to both membranes, although the percentages were intermediate between values for fresh and killed sperm. There was a tendency for a slightly greater percentage of the cryopreserved rooster sperm to bind to turkey membrane preparations than to those from chicken eggs. These results substantiate that capability of this general assay to provide data in which the percentage of sperm bound may be predictive of fertility does not require use of native vitelline membranes, but can be performed using protein extracts and, by logical extension, pure native protein(s) or the appropriate protein produced by recombinant technology and *in vitro* or *in vivo* expression of appropriate constructs.

Example 9

We took a microtiter plate and placed 110 μ L of the protein extracted from vitelline membrane (a concentration of 2.5 mg/mL protein in 20 mM Tris-HCl buffer, pH 8.0) preheated to 37° C into each well and dried the protein under low heat and vacuum. Then we placed 100 μ L of incubation buffer in each well together with an appropriate number of test sperm and incubated the wells for 2.5-3.0 hours at 37° C in a humidity chamber. After incubation, we decanted the contents of all wells and rinsed out each well repeatedly with excess incubation buffer. We fixed the cells to the membrane by exposure to a mixture of 3 parts ethanol and one part acetic acid and then stained the sperm with 4,6-diamidino-2-phenylindole (1 μ g/mL in PBS buffer). The number of sperm bound per unit area was quantified by counting using a fluorescence microscope. Data analysis was conducted as described in the foregoing examples.

For example, when we evaluated ability of these data to discriminate between samples of rooster spermatozoa representing fresh semen from unique males, we were able to show that samples differing in capability to bind to the egg membrane are likely to differ in fertilizing capability. Ten sperm samples were tested for binding individually, and as a pooled sample containing equal parts of all samples. Whereas the pooled sample gave a uniform cell binding percentage of about 69%, individual samples gave the respective results of 68%, 58%, 49%, 34%, 95%, 64%, 17%, 78%, 67% and 22%.

Example 10

Semen was available from two bulls known to differ in fertility, despite similarity in the percentage of motile and morphologically normal sperm in their semen. Aliquots of fresh, unprocessed semen were diluted, evaluated for percent motile sperm and subaliquots containing 0 to 7×10^6 sperm were placed into wells in a prepared microtiter plate prepared according to Example 9. Approximately linear dose-response curves were obtained for each sample, and the slopes of the curves were different (see Figure 4). Few killed sperm bound to the assay plate. These data demonstrated that the present method is a meaningful fertility assay.

Example 11

Frozen semen was available from two stallions. Aliquots of these cryopreserved semen samples were thawed and diluted with TALP buffer. Sperm concentration was determined by cytometer count. Subaliquots containing 0 to 9×10^6 sperm were placed into wells in a microtiter plate prepared according to Example 9. Approximately linear dose-response curves were obtained for each sample (Figure 3).

Example 12

Sperm isolated from the cauda epididymis of mice were collected into TALP buffer, the sperm were enumerated by cytometer count, and aliquots of $1-10 \times 10^6$ sperm were placed into wells in a prepared microwell plate according to Example 9. An approximate linear dose-response curve was obtained (Figure 4).

Example 13

Three samples of cryopreserved human semen were available, but fertility of the samples was unknown. After thawing, each sample was deglycerolated by stepwise addition of 3 volumes of TALP buffer and divided into two aliquots. One was not treated further (unwashed), while the second was processed by gentle centrifugation (300 x G for 10 minutes) and the sperm were resuspended. Percentage of motile sperm in each sample was estimated. Sperm concentration was determined by cytometer and subaliquots containing 0 to 7×10^6 sperm were placed into wells in a prepared microtiter plate according to Example 9. Approximately linear dose-response curves were obtained for each sample, and the slopes of the curves were different (Figure 5). Very few killed sperm, in aliquots rapidly cooled to -196°C and then thawed, bound to the binding surface. Thus the assay is useful in determining the utility of human sperm.

Although the invention has been described with reference to specific techniques and results above, the invention is only to be claimed insofar as is set forth in the accompanying claims.

WE CLAIM:

1. A method for determining the potential fertility of a sample of spermatozoa comprising the steps of contacting spermatozoa from a sample onto turkey egg membrane protein or chicken egg membrane protein, quantifying the binding which takes place between the spermatozoa and said protein, and directly and linearly correlating the binding thus quantified to yield a numeric assay of fertilizing potential.
5
2. The method according to claim 1 wherein said membrane protein is a vitelline membrane protein.
3. The method according to claim 1 wherein the step of contacting spermatozoa from a sample with turkey egg membrane protein or chicken egg membrane protein further comprises the steps of selecting either chicken eggs or turkey eggs, dissecting the vitelline membranes from the eggs thus chosen, washing said vitelline membranes, extracting protein from said vitelline membranes, coating a solid substrate with said protein, forming wells adjacent the protein coating thus formed and contacting said spermatozoa onto said protein coating.
5
4. The method according to claim 3 wherein said solid substrate is a microscope slide.
5. The method according to claim 3 wherein the step of quantifying the binding which takes place between the spermatozoa and said protein further comprises the step of visually examining said wells under microscopy.
6. The method according to claim 3 wherein the step of quantifying the binding which takes place between the spermatozoa and said protein further comprises the step of visually examining said wells under fluorescent microscopy.

7. The method according to claim 2 wherein said vitelline membrane protein is divided and solubilized into proteins being retained by a 10 kDa filter.

8. The method according to claim 3 wherein the washing is carried out with a cleansing buffer comprising 50 mM sodium phosphate and 145 mM sodium chloride, pH 7.2.

9. The method according to claim 3 wherein the protein is coated onto the solid substrate by dipping, spraying or rolling a solution containing 2.5 mg/mL protein in 20 mM Tris-HCl buffer, pH 8.

10. The method according to claim 3 wherein said wells are formed by overlaying a silicone rubber template atop the protein coating.

11. A device for assaying the potential fertility of spermatozoa comprising a solid support having coated thereon protein extracted from the vitelline membranes of chicken or turkey eggs.

12. A device according to claim 11 wherein said solid support is a microscope slide and further wherein a template overlaying said protein has holes therein to form reaction wells adjacent said protein.

13. A device according to claim 12 wherein said protein is 10 kDa in size.

14. A device for assaying the potential fertility of spermatozoa comprising a solid support having one or more preformed wells into which is coated extracts from the vitelline membrane of turkey or chicken eggs.

15. A device according to claim 14 wherein said extract is a protein having a size of 10 kDa.

16. A device for assaying the potential fertility of spermatozoa comprising a support bearing at least one vitelline membrane from a chicken or turkey egg.

17. The device according to claim 16 wherein said support is a solid support and at least one well is positioned adjacent said vitelline membrane.




FIG. 1




FIG. 2




FIG. 3




FIG. 4

2/3

SUBSTITUTE SHEET (RULE 26)




FIG. 5

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US95/05117

A. CLASSIFICATION OF SUBJECT MATTER

IPC(6) : C12M 3/04, 3/10; C12Q 1/02, 1/24

US CL : Please See Extra Sheet.

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 435/1, 4, 284, 287, 299, 301; 514/2; 424/520, 522, 588, 559, 561, 581, 582

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

APS, BIOSIS

search terms: vitelline, sperm, fertility

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X --- Y	Journal of Reproductive Fertility, Volume 80, issued 1987, G.J. Wishart, "Regulation of the length of the fertile period in the domestic fowl by numbers of oviducal spermatozoa, as reflected by those trapped in laid eggs", pages 493-498, especially page 494, paragraph 1.	1, 2 ---- 3-17



Further documents are listed in the continuation of Box C.



See patent family annex.

* Special categories of cited documents:	*T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
A document defining the general state of the art which is not considered to be of particular relevance	*X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
E earlier document published on or after the international filing date	*Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	*Z* document member of the same patent family
O document referring to an oral disclosure, use, exhibition or other means	
P document published prior to the international filing date but later than the priority date claimed	

Date of the actual completion of the international search

26 MAY 1995

Date of mailing of the international search report

08 JUN 1995

Name and mailing address of the ISA/US
Commissioner of Patents and Trademarks
Box PCT
Washington, D.C. 20231

Facsimile No. (703) 305-3230

Authorized officer

JANE WILLIAMS

Telephone No. (703) 308-0196

Form PCT/ISA/210 (second sheet)(July 1992)*

INTERNATIONAL SEARCH REPORT

International Application No.
PCT/US95/05117

A. CLASSIFICATION OF SUBJECT MATTER:
US CL :

435/1, 4, 284, 287, 299, 301; 514/2; 424/520, 522, 588, 559, 561, 581, 582



11